



Intravenous Thrombolysis in Stroke Patient with Previous ICH History and Warfarin Consumption Even for Second Time

Abdolreza Ghoreishi¹, Zeinab Golmohamadpour²

¹ Department of Neurology, Vali-e-Asr-Hospital, School of Medicine, Zanjan University of Medical Science, Iran

² General Practitioner, Zanjan University of Medical Science, Iran

Article Info	ABSTRACT
Article type: Case report	Background and Aim: According to the contraindication usage of IV thrombolysis for patients with previous ICH in guideline and Because of different outcome about use of IV TPA in patient with previous ICH & Warfarin consumption which hasn't reported yet, it made us try it And it had therapeutic result.
Article History: Received: 20 January 2020 Revised: 04 March 2020 Accepted: 14 May 2020	Case Report: Y/O woman with sudden left hemiplegia that she had ICH 6 months ago and because of AF was on warfarin. she was diagnosed with ischemic CVA and NIHSS:20 at first but after twice injection of IV TPA, finally discharged with NIHSS :3.
Keywords: Hemiplegia Ischemic CVA NIHSS Second injection TPA Warfarin	Conclusion: In this treatment procedure there was a CVA patient on warfarin consumption elected for IV thrombolysis and some complications happened :such as second CVA in one week later after first IV TPA injection, and then she had second IV thrombolysis but finally it caused to discharge symptom free. According to severity of CVA in any patient and past history of them, it's possible to use IV TPA injection.
